

## ПРИРОДНО - МАТЕМАТИЧКИ ФАКУЛТЕТ НИШ

## **KEY WORDS DOCUMENTATION**

Accession number, AN	IO:	
Identification number, I		<u> </u>
Document type, <b>DT</b> :		monograph
Type of record, <b>TR</b> :		textual / graphic
Contents code, CC:		university degree thesis
Author, <b>AU</b> :		Marko Rančić
Mentor, MN:		Ivan Mančev
Title, <b>TI</b> :		THE MOST FREQUENTLY USED BASIS SET OF FUNCTIONS IN THE MOMENTUM SPACE
Language of text, <b>LT</b> :		Serbian
Language of abstract, <b>LA</b> :		English
Country of publication, <b>CP</b> :		Serbia
Locality of publication, <b>LP</b> :		Serbia
Publication year, <b>PY</b> :		2012
Publisher, <b>PB</b> :		author's reprint
Publication place, <b>PP</b> :		Niš, Višegradska 33.
Physical description, PD: (chapters/pages/ref./tables/pictures/graphs/appendixes)		43 p. ; graphic representations
Scientific field, <b>SF</b> :		Physics
Scientific discipline, <b>SD</b> :		Atomic and molecular physics
Subject/Key words, <b>S/KW</b> :		Fourier transform, Slater orbital, Gauss orbital, hydrogen wave function, Coulomb Sturmian
UC		539.1
Holding data, <b>HD</b> :		Library
Note, <b>N</b> :		
Abstract, <b>AB</b> :		The wave function in momentum space is defined as the Fourier transform of the ordinary wave function in position space. In the present master work the Fourier transforms are carried out for the most frequently used basis set of functions, such as hydrogenic wave functions, Slater orbitals, Gauss orbitals as well as Sturmian orbitals.
Accepted by the Scientific Board on, ASB:		11.9.2012.
Defended on, <b>DE</b> :		
Defended Board, <b>DB</b> :	President:	
	Member:	
	Member, Mentor:	